


Seabird bycatch reduction in scampi trawl fisheries


JOHANNA P. PIERRE
JOHN CLEAL
FINLAY N. THOMPSON
EDWARD R. ABRAHAM

Draft final report prepared as
part of Department of Conservation project
MIT2011-02



www.dragonfly.co.nz

Dragonfly Science
PO Box 27535
Wellington 6141
New Zealand



www.fishinfo.co.nz

Clement & Associates Limited
Postal Box 1460
Nelson
New Zealand

To be cited as:

Pierre, J. P.; Cleal, J.; Thompson, F.N.; Abraham, E.R. (2013). Seabird bycatch reduction in scampi trawl fisheries. Final Research Report for Department of Conservation project MIT2011-02 (Unpublished report held by Department of Conservation, Wellington). 26 pages.



Crown copyright © Following the New Zealand Government Open Access and Licensing (NZGOAL) framework, this report is licensed under a Creative Commons Attribution 3.0 New Zealand Licence. This allows you to distribute, use, and build upon this work, provided credit is given to the original source.

EXECUTIVE SUMMARY

Seabird bycatch rates in the scampi fishery are estimated to be the second highest amongst New Zealand trawl fisheries. Seabird captures in this fishery are exacerbated by characteristics of the fishing operation: the gear is at or near the sea-surface for extended periods during shooting and hauling, and the catch typically comprises over 80% fish and invertebrate bycatch, which is discharged at the fishing grounds. A substantial body of work exists on seabird bycatch reduction measures for trawl fisheries. However, characteristics of scampi trawl gear and the prevalence of net captures amongst bycaught birds (for which no deployment-ready mitigation measures are available) present challenges for reducing seabird catch in this fishery. This project sought to identify potential methods with which to mitigate seabird captures in the New Zealand scampi fishery, test the feasibility and effectiveness of these methods, and make recommendations on future work on seabird bycatch in this fishery.

Through reviewing available information and holding an expert workshop (including representatives from the scampi fishing industry), we identified three areas for work: improving batch discharge regimes to ensure discharge is held on board during shooting and hauling; improving the design and construction of paired streamer lines; and testing the 'net restrictor' – a novel approach to reducing seabird captures in scampi nets. The first two areas of work will be addressed on an ongoing basis through working with skippers and crews, and utilising observer coverage of scampi vessels.

Deployment of the net restrictor prevents the mouth of the net from becoming wide open during shooting and hauling. First, we examined the operational feasibility of the restrictor in the centre net of a triple-rig targetting scampi. Then, we designed an experiment to test the efficacy of the restrictor in reducing seabird catch. Constraints on government observer coverage prevented the implementation of this experiment during the course of the project. However, implementing data collection protocols in future years on observed trips where vessels are using net restrictors will allow the assessment of the efficacy of the net restrictor in reducing seabird catch.

Footage collected using underwater cameras confirmed that the height of the centre net in triple-rig scampi gear was reduced by approximately 75% when restrictor ropes were in place. Footage also showed that the headline and some of the body of mesh around the headline sat lower in the water column with restrictors in place than without. While not a substitute for a designed experiment, this footage is a preliminary indication that the restrictor may be effective in reducing the risk of seabird bycatch in centre nets at shooting and hauling. We recommend empirical testing of the efficacy of net restrictors in this, and potentially other, demersal fisheries in which seabirds are caught in trawl nets.

1. INTRODUCTION

Scampi (*Metanephrops challengeri*) is a small, benthic crustacean fished by demersal trawling in New Zealand waters. In 2012–13, the total allowable commercial catch was 1190 tons. Annually, the scampi fishery produces 600–750 tonnes of product. The majority of the catch is taken from Quota Management Areas SCI1, SCI2, SCI3 and SCI6A, in the Bay of Plenty, Wairarapa Coast, Chatham Rise and around the Auckland Islands respectively (Ministry for Primary Industries 2012). Scampi typically represents less than 20% of the catch landed in the double- or triple-rigged nets deployed. Fish and invertebrate bycatch make up the remainder of the catch. Vessels normally process their catch at sea, and discharge unwanted bycatch at the fishing grounds (Ministry for Primary Industries 2012).

Seabird captures in the scampi fishery are exacerbated by a number of characteristics of fishing operations. The gear, consisting of multiple adjacent nets, can be at or close to the water surface for extended periods during shooting and hauling. Further, the netted catch and discharge of non-target bycatch from each haul attracts birds to vessels. Estimated total seabird captures for tows targeting scampi ranged from 67 to 443 birds annually for the fishing years 1998–99 to 2010–11 (95% c.i., Abraham et al. 2013). Observed capture rates ranged from 1.04 to 20.34 birds per 100 tows over the 1998–99 to 2011–12 period (Abraham & Thompson 2012). The average observed capture rate of seabirds in this fishery for the years 2002–03 to 2011–12 was 5.10 birds per 100 tows, which is the second highest seabird capture rate for New Zealand trawl fisheries. Seabird species that have been reported caught have included southern Buller's albatross (*Thalassarche bulleri*), Salvin's albatross (*Thalassarche salvini*), Campbell albatross (*Thalassarche impavida*), white-capped albatross (*Thalassarche steadi*), Cape petrel (*Daption capense*), black petrel (*Procellaria parkinsoni*), giant petrel (*Macronectes* spp.), white-chinned petrel (*Procellaria aequinoctialis*), flesh-footed shearwater (*Puffinus carneipes*), and sooty shearwater (*Puffinus griseus*) (Abraham et al. 2013). Of these species black petrel, flesh-footed shearwater, Salvin's albatross, and New Zealand white-capped albatross have been identified as species that are at risk of population-level impacts as a result of fatalities in New Zealand commercial fisheries (Richard & Abraham 2013).

A substantial body of research exists on methods to reduce seabird bycatch in trawl fisheries. Nets, warp cables, and monitoring cables (e.g., third wires and paravanes) are the components of fishing gear that may capture or kill seabirds during trawl fishing (Bull 2007, 2009, Løkkeborg 2011). However, despite common elements, the form and configuration of gear components differs between conventional finfish trawl fisheries and shrimp trawl fisheries. While seabird bycatch has been recorded in shrimp fisheries internationally (González-Zevallos et al. 2011, Marinao & Yorio 2011), mitigation measures have not been deployed to reduce this. Similarly, in New Zealand, scampi fishing involves gear and operational practices that are unlike other trawl fisheries, and mitigation approaches are not well developed as yet.

Due to the high seabird bycatch rates observed in the scampi trawl fishery, this project (Conservation Services Programme project MIT2011–02) aimed to develop methods to mitigate the capture of seabirds in this fishery. The specific objectives were:

1. To identify methods to mitigate the capture of seabirds in the commercial scampi trawl fishery,
2. To test the feasibility, and to the extent possible the effectiveness, of methods to mitigate the capture of seabirds in the commercial scampi trawl fishery, and,
3. To make recommendations for future work to develop and/or test the effectiveness of methods to mitigate the capture of seabirds in the commercial scampi trawl fishery.

2. METHODS

2.1 Methods to mitigate seabird captures in trawl fisheries

Our approach to identifying measures that may reduce seabird captures during scampi fishing operations was threefold:

- conduct a review of previous research on seabird bycatch reduction measures in trawl fisheries,
- review existing data collected by New Zealand government fisheries observers deployed in scampi fisheries and reported by fishers, and,
- hold a workshop with scampi fishers, scientists, fishery managers, and fisheries observers.

2.1.1 Review of mitigation measures

Measures to reduce the incidental capture of seabirds in trawl gear have focussed on finfish and squid fisheries to date (Bull 2007, 2009, Løkkeborg 2011). However, the attraction of seabirds to shrimp fisheries has been documented (Wickliffe & Jodice 2010) and seabird mortalities have been reported in shrimp trawl nets internationally (González-Zevallos et al. 2011, Marinao & Yorio 2011).

Consistent with shrimp fisheries in other locales (e.g., North and South America, see previous references), gear configuration used in the New Zealand scampi fishery is very different to trawl fisheries targeting finfish and squid. However, the same two gear components present the risk to seabirds in the scampi and other trawl fisheries, i.e., trawl warps and the trawl net. In New Zealand, seabird captures have been reported from both (see below for a detailed description of reported captures). In reviewing past research on methods intended to reduce seabird catch in trawl operations, we considered both warp and net captures. However, our focus was on net captures, given the relative prevalence of these during scampi fishing (Abraham et al. 2013, Pierre et al. 2012b). Further, while not enumerated as captures, another source of fatal interactions is seabirds colliding with fishing vessels (Rowe 2010, Ramm 2012).

There are three recent comprehensive reviews of measures designed to reduce seabird bycatch in trawl fisheries (Bull 2007, 2009, Løkkeborg 2011). Here, we consider these reviews and additional published reports to identify mitigation measures that might effectively reduce seabird bycatch on trawl warps and in trawl nets in the New Zealand scampi fishery.

Warp strikes

Currently, paired streamer lines (PSLs) are the international best practice device for reducing warp strikes. Streamer lines are also known as tori lines. Other measures have been investigated as possible mitigants of warp strikes, for example, bird bafflers, warp scarers, cones on warp cables and warp booms (summarised in Agreement on the Conservation of Albatrosses and Petrels (ACAP) (2011), and reviewed in detail in Bull (2007, 2009)). These measures have not been effective in consistently delivering reductions in warp strikes.

The efficacy of PSLs in reducing warp strikes has been demonstrated in a variety of fisheries, including off New Zealand (Middleton & Abraham 2007), the Falkland Islands (Reid & Edwards 2005, Sullivan et al. 2006a), and in the east Bering Sea (Melvin et al. 2011). The basic approach for trawlers is to attach one line to both the port and starboard stern quarters of a vessel, above and outside of the trawl blocks. Lines must have sufficient length and aerial extent to protect the full length of the trawl warp. Streamers

should be long enough to reach the water surface, and ideally brightly coloured. While PSLs are the best practice measure for reducing warp strikes, their effectiveness can be reduced in strong cross-winds. In such situations, the lines may not effectively track the location of the warps, and so seabird exposure to the warps can increase. A towed object at the end of the lines reduces this issue, but does not eliminate it (Sullivan et al. 2006b). Further, seabirds can strike the PSLs themselves (Middleton & Abraham 2007).

Net captures

Measures for reducing seabird catches in trawl nets are much less developed than those reducing warp strikes. Currently, no methods have been widely tested and demonstrated to be effective in reducing net captures on shooting and hauling trawl nets. However, net binding and net weighting are methods which may reduce seabird captures during net shooting (Hooper et al. 2003, Sullivan et al. 2004, Clement and Associates 2009).

Two operational measures are widely, and intuitively, considered to reduce the risk of net captures. These are minimising the period of time over which the trawl net is on the water surface, and, removing stuck fish and fish parts from the net prior to shooting (Hooper et al. 2003). The efficacy of these measures has not been tested quantitatively. However, the intuitive appeal of these measures has led to them being incorporated in Conservation Measures for the CAMLR Convention Area (through CCAMLR Conservation Measure 25-03), which are widely recognised as international best practice for seabird bycatch reduction.

Offal and discard management

Physically keeping birds away from potentially injurious fishing gear is desirable, for example, as addressed by PSLs above. However, the underlying reason seabirds attend vessels is due to the attraction of food, i.e. the trawled catch and discharged processing waste. Offal and discard management is a critical part of best practice for reducing seabird bycatch, which works by reducing the abundance of seabirds around vessels, consequently reducing the likelihood of captures.

Pierre et al. (2012a) summarises experimentally tested measures for offal and discard management. The preferred waste management option, i.e., the most effective in reducing bycatch risk, is to hold all processing waste for discharge when trawl gear is out of the water. If waste cannot be retained for the duration of trawl tows, discharging waste as quickly as possible (in maximally large batches and as infrequently as possible) is the recommended practice for reduction of seabird interactions with trawl warps. Holding waste for periods as short as 30 min may reduce the numbers of petrels and shearwaters (excluding Cape petrels) attending vessels. However, longer holding periods of up to 8 h may be required. Holding waste for periods of 2 h can reduce vessel attendance by albatrosses and giant petrels. However, longer holding periods (e.g., 4 h) may be required. Eight hour holding periods are preferable to 4 h holding periods, to further reduce seabird abundance at vessels.

Mincing processing waste can reduce the abundance of some seabirds at vessels. Mincing is most effective in reducing vessel attendance by great albatrosses (*Diomedea* spp.). It is less effective in reducing attendance by smaller albatrosses (e.g., *Thalassarche* spp.). Further, limitations of mincing machinery mean that not all processing waste can be minced and retrofitting vessels with mincers is costly.

Discharging waste as it becomes available during processing is the least desirable waste management option.

Not discharging fish processing waste before shooting or hauling of the net is another intuitive element of best practice which is considered to reduce both seabird captures both on trawl warps and in nets. This is because seabirds are less likely to be around the vessel (or will be in lower abundances around the vessel) in the absence of discharge. Consequently, they will not be available to be caught on trawl gear. The contents of nets will always be attractive to seabirds and may draw them to a haul event. However, even if some of the haul occurs prior to seabirds arriving in an area, risk of seabird captures is lower for that period.

Vessel strikes

While not a focus of this project, vessel strikes result in some seabird fatalities in the scampi fishery (Rowe 2010, Ramm 2012). Vessel strikes can result from different factors, e.g., extreme weather conditions (which may drive birds into vessels through wind gusts), birds becoming confused in deck lights, etc.. Across fisheries, best practice to manage this issue is to use deck lighting such that crew safety is ensured but seabird risk is minimised (Black 2005).

2.1.2 Review of existing data collected by New Zealand government fisheries observers and reported by fishers

Information extracted from Ministry for Primary Industries databases was explored in order to characterise the scampi fishery, including fishing effort, vessels involved, and seabird captures. Fisher-reported seabird captures were also examined (Pierre et al. 2012b). Seabird capture data described catch rates per unit of fishing effort, as well as seasonality of captures and gear factors potentially increasing capture risk. Observer comments were also considered, in identifying risk factors that may have contributed to bycatch events.

Fishing capacity and effort distribution in space and time

Fishing vessel and effort data were extracted from the Ministry for Primary Industries' Warehouse database. Since the 2000–01 fishing year, 8 - 19 vessels have targeted scampi in quota management areas (QMAs) SCI1 - SCI9 (Table 1). There was no fishing in SCI8 or SCI10, where TACCs have been 5 t and 0 t, respectively since 2003–04 (Clement and Associates 2011). Scampi was introduced into the Quota Management System (QMS) on 1 October 2004. Prior to that date, a catch limit existed (Clement and Associates 2011). Also on 1 October 2004, Quota Management Areas were given their current boundaries. Specifically, the boundary was shifted between SCI3 and SCI4A, and the area of SCI6A was adjusted from a circular to a trapezoid configuration. After introduction into the QMS, TACCs and reported catch did not change greatly (Clement and Associates 2011) although the fleet almost halved in size between the 2003–04 and 2004–05 fishing years (Table 1). In recent years, the number of vessels targeting scampi has been approximately proportional to each QMA's total allowable commercial catch (TACC), i.e., most vessels fished where TACCs were highest (SCI3 and SCI6A).

Approximately 4000 to 5000 trawl tows targeting scampi are conducted annually (Figure 1). Since introduction into the QMS, annual effort (number of trawl tows) has been most variable in SCI3 and SCI4A, and most consistent in SCI1 and SCI6A. Most fishing effort occurred between October and January, in QMAs SCI3, SCI1, and SCI2 (Figure 2). Least effort occurred between April and September. On average, the number of trawl tows targeting scampi in October was almost double that conducted in July (Figure 2).

Table 1: Fishing vessels targeting scampi, by year (1 October - 30 September) and area. 'Total' = all vessels fishing in the New Zealand EEZ each year.

	Quota management areas									Total
	SCI1	SCI2	SCI3	SCI4A	SCI5	SCI6A	SCI6B	SCI7	SCI9	
2000–01	9	9	10		2	10				12
2001–02	10	14	12	1		10	2	2		15
2002–03	9	12	16	6	4	9	2	2		19
2003–04	7	7	15	4	2	10				17
2004–05	5	8	8	8	2	6	1	1	1	9
2005–06	3	8	8	5	1	6	1	1		9
2006–07	3	6	7	5		6				10
2007–08	4	6	7	2	1	7				11
2008–09	3	5	9		1	5				9
2009–10	5	6	6	1		6	1			8
2010–11	3	4	7	4		7				8




Figure 1: Annual fishing effort (number of trawl tows targeting scampi) from the 2000–01 to the 2010–11 fishing years in Quota Management Areas. Colours represent Quota Management Areas, as shown in the legend. Effort is delineated according to current QMA boundaries.




Figure 2: Fishing effort targeting scampi (tows per year) by month, for the fishing years 2000–01 to 2010–11. Colours represent Quota Management Areas, as shown in the legend. Effort is delineated according to current QMA boundaries.

Table 2: Fisher-reported seabird captures in the scampi fishery, for fishing years 2001-2011. Codes are as used on the Non-fish Protected Species Catch Return.

Species	Code	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	Total
Albatrosses	XAL	3	3	5		20	16	15	22	13	14	15	129
Antipodean and Gibson's albatrosses	XAG										1		1
Black-browed albatrosses	XKM										1	1	2
Campbell black-browed albatross	XCM							1	4	2			2
Salvin's albatross	XSA									1		4	10
Southern Buller's albatross	XBM						3						3
Buller's albatrosses	XPB									7		7	14
White-capped albatrosses	XSY					2	2	2		1	8		13
New Zealand white-capped albatross	XWM									2	6		8
Petrels	XPE		1		2	11	1	9	4				28
Black petrel	XBP									2	3	2	7
White-chinned petrel	XWC							1				8	9
Flesh-footed shearwater	XFS								3	5	7	2	17
Common diving petrel	XDP										1		1
Sooty shearwater	XSH						5	3	13		5	32	58
Petrels, prions, and shearwaters	XXP									13	12	33	58
Seabirds	XSB										2		5
Small seabirds	XSS		2	1	3	14	13	9	1				50
Large seabirds	XSL		6	3	5	22	16	8	6	1			75
Total		11	11	13	11	67	56	48	53	49	58	104	481
Reported capture rate, %		0.22	0.16	0.25	0.29	1.44	1.15	0.93	1.10	1.23	1.37	2.34	0.91

Table 3: Seabird captures reported from the scampi fishery by government fisheries observers, percentage of trawl effort observed, capture rates per 100 tows, and estimated total captures (mean estimated captures with 95% confidence intervals). Data from Abraham et al. (2013)

	Effort	Observed			Est. captures	
		% obs.	Cap.	Rate	Mean	95% c.i.
2000/01	4978	5.3	9	3.38	139	104–178
2001/02	6719	8.8	6	1.02	167	123–216
2002/03	5130	9.98	8	1.56	158	93–271
2003/04	3753	10.98	8	1.94	114	67–191
2004/05	4648	3.08	9	6.29	212	135–330
2005/06	4867	6.8	13	3.93	225	142–351
2006/07	5135	7.58	24	6.17	157	106–227
2007/08	4804	10.91	11	2.1	163	102–251
2008/09	3975	9.96	19	4.8	209	137–316
2009/10	4248	8.19	5	1.44	180	112–285
2010/11	4447	12.05	86	16.04	307	225–433

Fisher-reported seabird captures

Under the Fisheries Act (1996), catches of protected species in commercial fisheries must be reported. Prior to 1 October 2008, fishers could use the Non-fish Incidental Catch Return form to report captures. These data are stored at NIWA (under contract to the Ministry for Primary Industries). After 1 October 2008, this form was discontinued and the Non-fish / Protected Species Catch Return form was deployed. Data from these forms are stored in the Warehouse database. Table 2 summarises seabird captures reported by fishers from 2001 to 2011, as extracted from the databases capturing data from these two forms.

The seabird capture rate reported by fishers has increased ten-fold from 0.22 in 2000–01, to 2.34 in 2010–11 (Table 2). Although greatly increased, it is still well below the capture rate reported by observers. For example, in 2010–11 observers reported a capture rate of 16.04 birds per 100 tows (Table 3). The range of species reported by fishers has increased greatly, with 9 different codes used in 2010–11, compared with three in 2000–01.

Observer-reported seabird captures

Government observers deployed in scampi fisheries also report seabird captures. Observer coverage rates have been relatively low in most years. However, capture events (Table 3) and potential risk factors associated with them (see sections below) have been documented. Modelling methods are used to develop fleet-wide estimates of seabird captures from these raw data (Table 3; Abraham et al. (2013)). Approximately three to twelve percent of trawl tows have been monitored by observers annually since the 2000–01 fishing year. Estimated total annual captures range from 114 to 307 on average. Confidence intervals (95%) vary from 67 to 433 captures (Table 3), highlighting the effects of low levels of observer coverage.

Table 4: Species of seabirds reported captured in the scampi fishery by government fisheries observers.

Fishing year	Species captured
2000–01	New Zealand white-capped albatross, Salvin’s albatross, Seabirds, white-capped albatrosses
2001–02	Black-browed albatross, Cape petrels, New Zealand white-capped albatross, Salvin’s albatross
2002–03	Albatrosses, common diving petrel, New Zealand white-capped albatross, Salvin’s albatross, sooty shearwater
2003–04	Albatrosses, Cape Petrel, New Zealand white-capped albatross, Salvin’s albatross, sooty shearwater, white-chinned petrel
2004–05	Chatham Island albatross, New Zealand white-capped albatross, Salvin’s albatross, southern Buller’s albatross, white-chinned petrel
2005–06	Albatrosses, black-browed albatrosses, flesh-footed shearwater, New Zealand white-capped albatross, petrels, southern Buller’s albatross
2006–07	Albatrosses, flesh-footed shearwater, New Zealand white-capped albatross, northern giant petrel, petrels, sooty shearwater
2007–08	Black petrel, flesh-footed shearwater, Salvin’s albatross, sooty shearwater
2008–09	Albatrosses, Cape Petrel, Cape petrels, flesh-footed shearwater, New Zealand white-capped albatross, Salvin’s albatross, southern Buller’s albatross
2009–10	Campbell black-browed albatross, flesh-footed shearwater, New Zealand white-capped albatross
2010–11	Flesh-footed shearwater, New Zealand white-capped albatross, Salvin’s albatross, sooty shearwater, white-chinned petrel

Species captured have included albatrosses, petrels, and shearwaters (Table 4). Observers have reported captures both on the trawl warps and in the trawl net. The majority of captures (albatrosses and petrels) have been detected in the trawl net. Albatrosses have also been captured on trawl warps/doors (five reported instances since 2007). In addition, two albatrosses have been reported caught on mitigation devices deployed (DOC and MPI, unpubl.).

Observer data entered in COD (the Ministry for Primary Industries’ Centralised Observer Database) may under-represent the number of seabirds captured in this fishery. This is because when multiple capture events occur, individual records have not consistently been completed for individual seabirds. This has been rectified where identified in the data. The land-based identification of bycaught seabirds returned from sea (Bell 2012) has highlighted this issue. This identification work encompasses the birds returned for necropsy as well as photos provided by observers. Two occasions have been identified where multiple capture events (15 and 10 birds) were observed in scampi trawl fisheries. In both cases the observer reported captures using a single entry on the nonfish bycatch form, and all captures were of white-chinned petrels. Where possible, and particularly for older records, verification of capture details is warranted using the forms on which these events were originally reported.

Observer comments on factors relevant to seabird bycatch risks

Observer comments relating to seabird captures were focused on discharge management and the use of mitigation devices. In 2007–08, one observer reported particularly high bird activity at full moon, and that a crew member actively shielded the net from birds using a plastic spade. Collection of offal and discards for batch discharge occurred on some vessels and others were equipped with a tank or hopper for storage of discharge but did not use it. There was some discharging ad hoc during shooting and towing. Streamer lines, bafflers and a buoy on a rope were all reported as being used with the intent of reducing seabird interactions (Rowe 2010). Observers also reported that seabirds were consistently attracted to the net. Similar observations were reported in 2008–09 and 2009–10 fishing years (Ramm 2010, 2012).

These observations suggest that improving batch discharge practices would reduce the risk of seabird bycatch. There also seems to be potential for work improving the design and construction of mitigation measures in use (e.g., streamer lines).

Exploratory analysis of gear characteristics that may influence bycatch risk

There have been 78 observed trawl scampi trips, on 15 different vessels, during the fishing years 2000–01 to 2010–11. Approximately a third of observed trips have occurred on the east coast of the South Island (SCI3), 28% around the Auckland Islands (SCI6A), and the rest split between SCI1, SCI2, and SCI4A. There has been very little observer coverage in SCI5 or SCI7.

In the 2007–08 fishing year observers started reporting gear configuration in detail on the MPI Trawl Gear Details Form. Information collected on this form was used to explore gear characteristics that may influence bycatch risk. There are 19 trips on seven vessels in the four years 2007–08 to 2010–11 for which gear configuration data were reported. Of those 19 trips, all but two have reported the use of tori lines. These two trips without tori lines occurred in the SCI1 and SCI3 areas, in the 2010–11 fishing year.

A step analysis (Akaike 1974, Venables & Ripley 2002) was applied to the observer records where gear configuration data were collected (Table 5). This included 950 tows. A negative binomial link function was used.

Results of step analysis

When all bird captures were considered, the most important factors (of those listed in Table 5) were month and area of fishing, followed by the number of nets and presence of a tori line. Codend mesh size was of marginal significance (Table 6).

Month of the year accounted for most variation in the data. Significantly fewer captures than expected were reported in August, November, and December. Captures increased in October, but this effect was not significant. Fewer captures than expected were reported in SCI2, 3 and 6A. However, the conclusions here are particularly vulnerable to the timing and location of observer coverage.

Significantly more captures were reported when three nets were deployed, compared to two nets. Increasing codend mesh size was related to a trend (of marginal significance) of increased captures.

Use of paired streamer lines reduced total seabird captures significantly.

Fewer albatrosses were captured in gear comprised of three nets compared to two (coefficients: $P < 0.05$). Headline length was of marginal significance, with a trend for more seabird captures in gear with longer headlines (Table 7).

For small birds, month, area, number of nets and codend mesh size were all significant in accounting for captures. August and December showed significantly lower captures than expected. Also, fewer captures than expected were reported in SCI2, 3 and 6A. However, as above, the conclusions here are particularly vulnerable to the timing of observer coverage.

Significantly more captures occurred during deployments of three nets compared to two, and increased captures occurred with increasing mesh size.

Table 5: Factors used in the step analysis. Codes used follow Sanders & Fisher (2010). Gear events: Z - No events, F - Haul, turn, reshoot - during a tow the vessel partially hauls the net, completes a turn, then reshoots the net, A, B - Net damage - where observer reported torn nets and nets that came fast, G, I - Gear breakage - where observer reported twisted warps, crossed doors, warp breakages, and lost gear, D - Winch failure at set, O, Y - Other - includes where observer has listed more than three, U - Unknown; Door type: H - High aspect, L - Low aspect, O - Other; Headline tag: (describes the source of headline height information) 1 - from net sonde measurements, 2 - a standard figure e.g., taken from net plans, 3 - from skipper.

QMA	SCI1 - 9
Number of codends	1, 2 or 3
Gear events	Mostly Z, but also A, AB, AIB, AO, B, BA, BG, D, F, G, GO, O, U, Y
Door spread	Range from 24 to 105 m
Lengthener mesh size	Range from 50 to 115 mm
Door type	H, L, or O
Month	Calendar month of fishing
Headline length	18 to 150 m
Headline tag	1, 2, or 3
Codend mesh size	Range from 50 to 115 mm
Trawl wingless	Yes or No
Number of warps	1 or 2
Tori lines used	Yes or no
	(No bafflers were reported in use on the vessels used in this analysis)
Vessel key	One of 7 vessels

Table 6: Summary of step analysis of all bird captures in scampi tows conducted during 19 trips on seven vessels in fishing years 2007–08 to 2010–11.

	Degrees of freedom	Deviance		
		Residual	Explained	Explained (%)
		356.85		
Month	10	213.76	143.09	40.1
Quota management area	3	176.13	37.63	17.6
Number of codends	2	158.49	17.64	10.0
Tori line present	1	148.36	10.12	6.4
Codend mesh size	1	144.86	3.50	2.4

Table 7: Summary of step analysis of albatross captures in scampi tows conducted during 19 trips on seven vessels in fishing years 2007–08 to 2010–11.

	Degrees of freedom	Deviance		
		Residual	Explained	Explained (%)
		76.34		
Number of codends	2	72.16	4.18	5.5
Headline length	1	68.98	3.18	4.4

Table 8: Summary of step analysis of small bird captures in scampi tows conducted during 19 trips on seven vessels in fishing years 2007–08 to 2010–11.

	Degrees of freedom	Deviance		
		Residual	Explained	Explained (%)
		336.71		
Month	10	183.92	152.78	45.4
Quota management area	3	141.48	42.45	23.1
Number of codends	2	121.42	20.06	14.2
Codend mesh size	1	117.64	3.77	3.1

Conclusions

The results of the step analyses should be interpreted cautiously given the limited amount of data available overall, as well as spatial and temporal constraints on data collected. Despite this, the analysis identified factors relating to gear deployment that may affect captures, i.e., number of nets, use of tori lines, and codend mesh size. Of these, codend mesh size had not been previously identified by observers as a possible risk factor. However, it has been explored to a limited extent by mitigation practitioners as a possible correlate of seabird captures in other fisheries (pelagic icefish trawl (*Champscephalus gunnari*), Roe (2005)), and is not currently supported as an avenue of mitigation research (Agreement on the Conservation of Albatrosses and Petrels (ACAP) 2011).

We recommend that this analysis is repeated when observer records for future fishing years are available.

2.1.3 Expert workshop

While the observer-collected information that is available is of high quality, the limited coverage of the scampi fleet by government observers results in the collective understanding of this fishery being less developed than for most other trawl fisheries. To complement the information extracted from available databases, and to progress the project from desktop work to at-sea information collection and trials, we held a workshop involving vessel management, skippers, crew, scientists, fishery managers, and government observers. This workshop included discussion of mitigation measures currently in use, issues around current measures, best practice measures used in other trawl fisheries, and potential new measures that could be explored (Pierre et al. 2012b).

The objectives of the workshop were:

- to seek feedback on factors identified as increasing the risk of seabird bycatch in the scampi fishery;
- to identify possible solutions to the risk factors identified;
- to discuss potential mitigation measures identified, and,
- to identify key information for observers to collect, relating to fishing practices and events that increase seabird bycatch risk.

In addition to Dragonfly and Clement and Associates (J. Pierre and J. Cleal), workshop participants included the scampi fleet manager from Sanford Ltd, three vessel skippers, one current and one former

Ministry of Fisheries observer, and representatives from the Department of Conservation and the Ministry for Primary Industries.

The workshop covered the following areas:

- a characterisation of the scampi fleet,
- vessel operations,
- current practices in the fleet that relate to seabird bycatch reduction,
- nature and extent of seabird captures,
- observer comments,
- possible mitigation measures, and,
- next steps.

Discussion at the workshop was wide-ranging and constructive, and is summarised below.

Batch discharging

Skippers were supportive of batch discharging as an efficient and effective way to reduce seabird activity around vessels. All reported having robust batching systems on their vessels. There was some discussion about the ease of emptying storage tanks. A variety of systems exist on different vessels, including hydraulic lifts and opening holes in tanks. Sometimes waste can become packed in towards the bottom of tanks, or when going through smaller openings. It is then dislodged manually. With one tonne capacity, tanks would not typically be emptied more than twice per tow; a large shot could land about 4 tonnes. There was some discussion about birds knowing they should hang around until the offal bin was emptied. On one vessel, the tank cannot be tipped when shooting/hauling due to the hydraulic arrangement involved. However, on other vessels, discharge during shooting and hauling had been reported in recent years by observers. The group discussed ways to improve performance in that area, e.g., ensuring crew coordination of shooting/hauling and discharging.

Paired streamer lines

Several observer photos of tori lines deployed from scampi vessels show that the design of these lines could be improved to increase efficacy. For example, lines could be deployed from greater heights, and incorporate more streamers. Skippers reported no difficulties in using tori lines, but generally did not consider them effective in deterring seabirds from the stern of vessels. This could be a consequence of designs currently in use.

While warp strikes have been reported, these are currently a secondary consideration to net captures in this fishery due to the short length of warps that is exposed (compared to in other trawl fisheries). However, offal is discharged when trawl nets are in the water, which means the discharge stream will flow past warp(s) on occasion. Observers have noted warp strikes occurring in this situation.

While the risk of tangling would have to be considered carefully, one skipper suggested that deploying a tori line during hauling may be useful for keeping birds away from the net area. Floats could be attached to the tori line to maintain buoyancy and reduce tangling.

Net captures

Discussion focussed on the risk of seabird captures in the middle net in triple-rig gears. Under typical operations with a triple rig, the middle net is held open including when the other two nets are being hauled.

The group speculated on where and when birds might be getting caught in the net. This is uncertain, and the group concluded that some footage taken astern should be informative in this regard. Similarly, observers could stand in a safe place astern and gather more information about how birds and the codends are interacting. Skippers and observers considered that net captures were probably occurring both on the shot and haul. One possible way to reduce seabird access to the net would be to limit the size (vertical height, from groundrope to headline) of the opening when the net is astern. At fishing depth, the vertical height is estimated at 1.5 to 2 m, so providing for this height to be constrained when the gear is not fishing should not affect fishing efficiency. This method was dubbed 'net restrictor'. Another measure that might be effective in reducing albatross access is dragging the net under the surface when it is not at fishing depth.

Net cleaning was recognised as a valuable part of practice to reduce the risk of seabird bycatch in (net) fisheries. However, skippers did not consider it feasible for scampi fishing due to the particular gear arrangement.

The group discussed the issue of triple rigs specifically. These are used to increase fishing efficiency but were also considered to increase bycatch risk (specifically due to the way the centre net is used). The third net is simple to detach (however obviously fishing efficiency is lost in this case). In cases of high bird captures/risk, if the third net is taken off, a key issue is when to put it back on again.

General issues

Health and safety on the vessel is paramount, and participants recognised that the development of any mitigation measures required consideration of deck safety issues in particular.

Other mitigation ideas

A water spray set-up was suggested as potentially useful for dispersing particularly large aggregations of birds. However, the group recognised that birds would probably get used to this measure if deployed constantly. To maintain effectiveness, it would have to be deployed in an unpredictable way, and perhaps reserved e.g., for when bird aggregations were particularly large or foraging aggressively.

Reducing the visibility of food in the net could be another approach to reduce seabird activity astern the vessel. There was some discussion about the use of dye to achieve this in exceptional circumstances, for example, a burly cage with a dye block inside could be deployed astern.

3. RESULTS

3.0.4 Application of mitigation measures to the scampi fishery

Given the extent of knowledge on the efficacy of batch discharge in reducing seabird bycatch risk (Pierre et al. 2012a), and the presence of holding tanks on scampi vessels, ensuring effective discharge

management is recommended to minimise capture risk. Batch discharge approaches could be refined to ensure robust regimes are implemented continuously on vessels. This can be achieved through crew education. Discharge has occurred during shooting and hauling in the past. Addressing this operationally should be relatively straightforward given the pre-existing hardware in place, but requires ongoing diligence from skippers and crew. Future observer coverage could also be used to monitor discharge practices.

The deployment of well-designed PSLs as warp strike mitigation measures has potential in the scampi fishery. Key considerations will be design of the PSLs and the identification of safe and effective locales for deployment and retrieval of the lines. Designs of PSLs deployed must be targeted to scampi gear, i.e. warp length and location astern. The design of paired streamer lines is expected to improve following consideration of recommendations and outputs of Conservation Services Programme MIT2011–07 which examined the design of streamer lines used on trawl vessels ≥ 28 m in overall length (Conservation Services Programme 2011). Although this project focused on larger vessels, its recommendations provide for the development of streamer lines suitable for different sized vessels (Cleal et al. 2013).

Net capture mitigation measures are less simply transferred from other fisheries to scampi, and development of these measures is in its infancy across trawl fisheries. The net is typically not brought onboard in its entirety during scampi fishing. Nets can also be on the surface for relatively extended periods, as the rig is comprised of multiple nets. Net mouths being open and closing during the haul is expected to be a key source of bycatch risk. Measures used in other trawl fisheries such as net binding and weighting may be worth investigating further. The most promising avenue for further work was considered to be the net restrictor.

To test the feasibility of the net restrictor at sea, restrictor ropes were deployed by one vessel using a triple-rig over a period of several months. Restrictor ropes were fitted across the mouth of the centre trawl net (Figure 3). In that time, skippers and crew monitored the performance of the restrictor ropes. Ropes did not tangle, and were readily attached to the headline and groundrope of the net. While not evaluated quantitatively, crew considered that the handling of the centre net improved with restrictors fitted. In addition, while not tested empirically, crews felt that fish and invertebrate bycatch was lower in the net when the restrictor ropes were in place. Given the feasibility of the method from an operational perspective, moving to a trial of the restrictor's efficacy in reducing seabird bycatch was warranted.

3.1 At-sea testing of the net restrictor

3.1.1 Experimental design for testing the restrictor at sea

At the outset of the project, government observer coverage of scampi fishing activity was the platform expected to support testing of potential mitigation measures at sea, should new measures be identified (Conservation Services Programme 2011, Pierre et al. 2012b). Planned observer coverage for the year from July 1 2011 to June 30 2012 was 450 observer days (Conservation Services Programme 2011). Once the operational feasibility of the restrictor was confirmed, we designed an experimental approach using normal observer coverage to test the efficacy of the restrictor in reducing seabird catch in the centre net of triple-rig scampi trawl gear.

The experimental design specified the deployment of restrictors on the centre net of triple-rigged vessels, in accordance with a pre-defined randomised schedule. Restrictors were to be deployed randomly for blocks of two to four consecutive treatment days, with treatment days starting and ending at midnight. If operational reasons (e.g., bad weather or fishing occurring on muddy substrates) required the removal of




Figure 3: Photos from the at-sea feasibility trial of the net restrictor. (a) the restrictor (red rope) affixed to the net rope, (b) the restrictor spanning the mouth of the trawl net, (c) the restrictor in place in the mouth of the net, as the net is deployed.

the centre net, the treatment days were to be paused, and then resumed when fishing involving the centre net recommenced. For example, if muddy substrates were fished two days into a four day block when restrictors were to be deployed, the centre net was removed. After that fishing activity concluded, the centre net was to be redeployed with restrictors in place, and the second two days of the four day block were to be fished.

The restrictor arrangement consisted of six ropes that were to be fitted across the mouth of the centre net, i.e., from the headline to the groundrope. One restrictor rope was to be fitted 1 m to each side of the headline centre. Another 2 restrictor ropes were to be fitted to each side of the net mouth, 2 m apart (Figure 4). Restrictor ropes were made of 6–10 mm nylon braid and were either attached using non-slip knots, or using an alternative attachment method which facilitated their removal during the experiment. The length of restrictor ropes matched the headline height of the net, allowing the net to keep its intended shape during fishing. Observers were tasked with checking the placement of the restrictor ropes prior to each tow, and confirming that the experimental treatment days were followed in the random order prescribed prior to sailing.

In lieu of a non-lethal metric with which the efficacy of the restrictor at reducing seabird bycatch could be determined, the response variable for the experiment was to be seabird captures. As part of their




Figure 4: Experimental set-up of the net restrictors across the mouth of the centre net in triple nets used for scampi trawling.

normal duties, observers record seabird captures and associated metadata on a series of forms. An additional form was designed for this experiment (Figure 5), which recorded the net (port, starboard, centre) in which captures occurred, and whether birds were caught on the restrictor rope itself. Observers were also to complete three fields that linked the experimental form to their other forms which recorded data including gear specifications, and details of target catch and bycatch. Finally, as this was a new experimental protocol, observers were encouraged to document any additional observations relating to the protocol, the restrictors, interactions with seabirds, or the experiment in general.

Data collected were expected to be analysed using generalised linear models executed in a Bayesian statistical framework, to determine whether restrictors were effective in reducing the risk of seabird captures.

Due to limited capacity within the government observer programme throughout this project, only one observed trip occurred on a scampi vessel testing the net restrictor. However, this trip was successful in implementing the experimental approach, and so confirmed the feasibility of conducting the experiment in the course of a normal fishing trip. The skipper reported that inserting and removing restrictors repeatedly was cumbersome as crew were not able to spread out the centre net on the deck of the vessel to measure the distance between restrictors. Marking the points along the headline and groundrope of the trawl net at which the restrictors were to be attached would address this.

3.1.2 Camera deployment

To understand the effects of the net restrictor on gear configuration (thereby confirming whether or not the restrictor reduced the size of the net mouth underwater astern the vessel) we worked with the research crew conducting stock survey work in Quota Management Area SCI6A on the F.V. San Tongariro, to deploy net-mounted underwater cameras (Figure 6). Two test tows using a Hampidjan 50 m scampi trawl were conducted on 24 March 2013, in waters north of the Auckland Islands. Restrictors were not deployed during the first tow, and eight restrictor ropes were deployed across the centre net during the second tow (Figure 7). The gear was monitored with two GoPro cameras in place. These cameras were




Figure 6: Camera deployed in the centre net of a triple-rig scampi trawl.




Figure 7: Net hauling on the vessel from which tows were recorded during trials of net restrictors.

than the groundrope camera. This was due to a more secure fitting, and because the meshes behind the groundrope were less stable than behind the headline. Movement of the groundrope during towing resulted in the camera affixed there capturing a highly dynamic view such that it was often not clear which part of the net featured in the footage.

During filming, the centre net spread to approximately 15 m (estimated wing end spread). Without restrictors in place (Figure 9), the height of the centre net mouth was measured as 4–4.5 m. With restrictors in place (Figure 8), this diameter decreased to a maximum of 1.0–1.2 m at 48 m depth, with lower headline heights detected during shooting and hauling. In addition to the altered net dimensions that restrictors achieved, footage also showed that the headline and some of the body of mesh around the headline sat lower in the water column with restrictors in place than without. When restrictors were not in place, the trawl headline was clearly visible at the water surface and associated meshes floated as well.

(a)



(b)




Figure 8: Net restrictors across the central net of a triple-rig scampi trawl during hauling.




Figure 9: Deployed scampi trawl without net restrictors in place.

The weight of the groundrope (350–400 kg) was observed to pull the headline down below the water surface more rapidly when restrictors are in place.

4. DISCUSSION

Three main areas were identified for further work aimed at reducing seabird captures in the scampi fishery. These were the implementation of robust batch discharge regimes, improvement of streamer line designs, and the experimental deployment of the net restrictor. Ongoing crew education, combined with feedback from government fisheries observers monitoring vessels at sea will facilitate ongoing implementation of robust batch discharge regimes on scampi vessels. Recommendations have been developed for the design and construction of paired streamer lines on vessels ≥ 28 m. These recommendations have been promulgated amongst the deepwater trawl fleet in the form of a fact sheet (Cleal et al. 2013). While streamer lines described by these recommendations will require some tailoring to smaller scampi vessels, the principles of good design will be similar (e.g., ensuring aerial extent and warp coverage).

Constraints on the delivery of government observer services prevented the empirical testing of the net restrictor at sea. However, an experimental design and protocol were developed and trialled, with which this potential mitigation measure could be tested in future. Footage collected from the centre net of a triple-rig confirmed that the restrictor was effective in reducing the size of the net opening. The footage also showed that meshes around the headline sat lower in the water during shooting and hauling operations when restrictors were in place. Based on these characteristics, the restrictor is expected to be effective in reducing seabird bycatch risk created by the mouth of the trawl net and possibly also by reducing the lofting of meshes around the headline at the sea surface.

The deployment of net restrictors is not currently required in the scampi fishery. However, in lieu of a designed experiment, deploying the restrictor over time will facilitate determination of its efficacy, in terms of both the rate of seabird bycatch and the volume of fish and invertebrate bycatch landed. To this end, we recommend that for the next five years, the data collection form attached (Figure 5) is distributed to observers on scampi vessels that are using restrictors in the centre net of triple-rig gear, and that this form is completed together with regular forms describing seabird, fish and invertebrate bycatch. While not a controlled experiment, this level of data collection would allow a quantitative comparison of seabird (and fish and invertebrate) bycatch rates before and after the restrictors were deployed.

Currently, the scampi fishery has the second highest rate of observed seabird bycatch of any New Zealand trawl fishery (excluding inshore trawl fisheries, for which observer coverage in most areas is insufficient to support the robust estimation of bycatch rates). Seabirds reported captured from the scampi fishery include species of particular conservation concern, given that captures made in the course of commercial fishing exceed the species' sustainability limits (e.g., Salvin's albatross, flesh-footed shearwater, and black petrel, Richard et al. (2011), Richard & Abraham (2013)). While not a substitute for a designed experiment, preliminary results from camera footage show that the net restrictor may be effective in reducing the risk of seabird bycatch in centre nets. We recommend the empirical testing of the efficacy of net restrictors in this, and potentially other, demersal fisheries where seabirds are caught in trawl nets.

5. ACKNOWLEDGEMENTS

We appreciate the input, assistance and cooperation of the following skippers and crew, fleet, vessel and fishery managers, scientists, and observers in undertaking this work: M. Beritzhoff-Law, H. Butler, J. Clement, L. Duggan, J. Finlayson, S. Gibb, S. Hornby, D. Middleton, K. O'Brien, K. Ramm, and D. Shaw. S. Gibb proposed the net restrictor concept. We are grateful to Dr I. Tuck and the crew of the F.V. San Tongariro conducting the SCI6A stock survey for managing the collection of footage of the net restrictor, with support from the National Institute of Water and Atmospheric Research.

This work was funded by the Department of Conservation's Conservation Services Programme as project MIT2011-02, principally through a levy on the quota holders of relevant commercial fish stocks.

6. REFERENCES

- Abraham, E.R.; Thompson, F.N. (2012). Captures of birds in scampi trawl fisheries, in the new Zealand exclusive economic zone, from 2002 to 2011. Retrieved from <http://data.dragonfly.co.nz/psc/v20121101/birds/scampi-trawl/all-vessels/eez/all/>, Apr 15, 2013.
- Abraham, E.R.; Thompson, F.N.; Berkenbusch, K. (2013). Estimated capture of seabirds in New Zealand trawl and longline fisheries, 2002-03 to 2010-11. Final Research Report for project PRO2010/01 (Unpublished report held by Ministry for Primary Industries, Wellington).
- Agreement on the Conservation of Albatrosses and Petrels (ACAP). (2011). ACAP summary advice for reducing impact of pelagic and demersal trawl gear on seabirds. Agreement on the Conservation of Albatrosses and Petrels, Hobart.
- Akaike, H. (1974). A new look at the statistical model identification. *IEEE Transactions on Automatic Control* 19: 716-723.
- Bell, E. (2012). Identification of seabirds captured in NZ fisheries: 2011/12. Department of Conservation, New Zealand.
- Black, A. (2005). Light induced seabird mortality on vessels operating in the Southern Ocean: incidents and mitigation measures. *Antarctic Science* 17(01): 67-68.
- Bull, L.S. (2007). Reducing seabird bycatch in longline, trawl and gillnet fisheries. *Fish and Fisheries* 8: 31-56.
- Bull, L.S. (2009). New mitigation measures reducing seabird by-catch in trawl fisheries. *Fish and Fisheries* 10: 408-427.
- Cleal, J.; Pierre, J.P.; Clement, G. (2013). Warp strike mitigation devices in use on trawlers ≥ 28 m in length operating in New Zealand fisheries. Department of Conservation, Wellington, New Zealand. Available from <http://www.doc.govt.nz/publications/conservation/marine-and-coastal/conservation-services-programme/csp-reports/warp-strike-mitigation-devices/> 42 p.
- Clement and Associates. (2009). Mitigating seabird interactions with trawl nets: A report commissioned by Department of Conservation Project 2006-02. Department of Conservation, Nelson, New Zealand. Available from <http://www.doc.govt.nz/publications/conservation/marine-and-coastal/conservation-services-programme/csp-reports/archive/2008-2009/mit2006-02-final-report-mitigating-seabird-net-captures-in-trawl-fisheries/> 70 p.
- Clement and Associates. (2011). New Zealand commercial fisheries: The atlas of area codes and TACCs. 2011/2012. Clement and Associates Ltd., Nelson, New Zealand
- Conservation Services Programme. (2011). Marine Conservation Services Annual Plan 2011/12. Department of Conservation, Wellington. 70 p.
- González-Zevallos, D.; Yorio, P.; Svagelj, W.S. (2011). Seabird attendance and incidental mortality at shrimp fisheries in Golfo San Jorge, Argentina. *Marine Ecology Progress Series* 432: 125-135.
- Hooper, J.; Agnew, D.; Everson, I. (2003). Incidental mortality of birds on trawl vessels fishing for icefish in subarea 48.3. *CCAMLR WG-FSA-04-03*.

- Løkkeborg, S. (2011). Best practices to mitigate seabird bycatch in longline, trawl and gillnet fisheries – efficiency and practical applicability. *Marine Ecology Progress Series* 435: 285–303.
- Marinao, C.J.; Yorio, P. (2011). Fishery discards and incidental mortality of seabirds attending coastal shrimp trawlers at Isla Escondida, Patagonia, Argentina. *The Wilson Journal of Ornithology* 123(4): 709–719.
- Melvin, E.F.; Dietrich, K.S.; Fitzgerald, S.; Cardoso, T. (2011). Reducing seabird strikes with trawl cables in the pollock catcher-processor fleet in the eastern Bering Sea. *Polar Biology* 34: 215–226.
- Middleton, D.A.J.; Abraham, E.R. (2007). The efficacy of warp strike mitigation devices: Trials in the 2006 squid fishery. Final Research Report for research project IPA2006/02. (Unpublished report held by Ministry for Primary Industries, Wellington).
- Ministry for Primary Industries. (2012). Report from the Fisheries Assessment Plenary, May 2012: stock assessments and yield estimates. Ministry for Primary Industries, Wellington, New Zealand. 157 p.
- Pierre, J.P.; Abraham, E.R.; Richard, Y.; Cleal, J.; Middleton, D.A.J. (2012a). Controlling trawler waste discharge to reduce seabird mortality. *Fisheries Research* 131–133: 30–38.
- Pierre, J.; Cleal, J.; Thompson, F.N.; Butler, H.; Abraham, E.R. (2012b). Seabird mitigation in New Zealand’s scampi trawl fishery. Progress Report (Unpublished report held by Department of Conservation, Wellington).
- Ramm, K. (2010). Conservation Services Programme Observer Report, 1 July 2008 to 30 June 2009. Final Report, Department of Conservation, Wellington, New Zealand.
- Ramm, K. (2012). Conservation Services Programme Observer Report, 1 July 2009 to 30 June 2010. Final Report, Department of Conservation, Wellington, New Zealand.
- Reid, T.; Edwards, M. (2005). Consequences of the introduction of tori lines in relation to seabird mortality in the Falkland Islands trawl fishery, 2004/05. Falklands Conservation, Stanley.
- Richard, Y.; Abraham, E.R. (2013). Application of Potential Biological Removal methods to seabird populations. *New Zealand Aquatic Environment and Biodiversity Report No. 108*. 30 p.
- Richard, Y.; Abraham, E.R.; Filippi, D. (2011). Assessment of the risk to seabird populations from New Zealand commercial fisheries. Final Research Report for research projects IPA2009–19 and IPA2009–20. (Unpublished report held by Ministry for Primary Industries, Wellington).
- Roe, R. (2005). Report and recommendations on mitigation use in the pelagic icefish (*Champscephalus gunnari*) fishery. Falklands Conservation, Stanley.
- Rowe, S.J. (2010). Conservation Services Programme Observer Report, 1 July 2007 to 30 June 2008. *DOC Marine Conservation Services Series 4*. Department of Conservation, Wellington, New Zealand.
- Sanders, B.M.; Fisher, D.O. (2010). Database documentation for the Ministry of Fisheries Centralised Observer Database. *NIWA Fisheries Data Management Database Documentation Series*.
- Sullivan, B.J.; Brickle, P.; Reid, T.A.; Bone, D.G.; Middleton, D.A.J. (2006a). Mitigation of seabird mortality on factory trawlers: Trials of three devices to reduce warp cable strikes. *Polar Biology* 29: 745–753.
- Sullivan, B.; Liddle, G.; Munro, G. (2004). Mitigation trials to reduce seabird mortality in pelagic trawl fisheries (Subarea 48.3). *CCAMLR WG-FSA-04-80*.
- Sullivan, B.J.; Reid, T.A.; Bugoni, L. (2006b). Seabird mortality on factory trawlers in the Falkland Islands and beyond. *Biological Conservation* 131: 495–504.
- Venables, W.N.; Ripley, B.D. (2002). *Modern applied statistics with S* (4th. ed.). Springer, New York.
- Wickliffe, L.C.; Jodice, P.G.R. (2010). Seabird attendance at shrimp trawlers in nearshore waters of South Carolina. *Marine Ornithology* 38: 31–39.